

raised. The snake was restrained in a 1-litre measuring glass cylinder and opening around its body was covered with cotton. External examination revealed a depression at the mid-dorsum (Image 1^w). Hence a radiograph was taken to identify any spinal injury. Radiograph revealed fractured vertebrae (Image 2^w).

A two inch wide micro pore adhesive tape was applied to the body of the snake to a length of 10in including the injured segment. A 50ml disposable plastic syringe, cut at both the ends and slit longitudinally into two equal halves, made smooth by filing on hard surface, padded suitably with cotton and gauze, was placed as splint on the lateral aspect of the body over the micro-pore bandage previously applied (Image 3^w).

The volunteers were advised to keep the snake in a limited confinement. The snake was again presented after 20 days for removal of the bandage. After restraining the micro-pore tape was cut open (Image 4^w). On releasing the snake on the ground it exhibited slow movement. The volunteers were advised to keep it in confinement for another 15 days. The snake recovered uneventfully. The volunteers reported that they have released the snake in its natural habitat.

^w See Images 1-4 in the web supplement at www.zoosprint.org



VET BRIEF

ZOOS' PRINT JOURNAL 22(6): 2736

Bone plating for tibial fracture repair in a Black Buck *Antelope cervicapra*

R.V. Suresh Kumar¹, S. Bharathi², G. Kamalakar³, P. Veena⁴ and Thoiba Singh⁵

^{1,2,4} Assistant Professors, Surgery and Radiology; ³P.G. Student, College of Veterinary Science, Tirupathi, Andhra Pradesh 517502, India
⁵ Veterinary Assistant Surgeon, S.V. Zoological Park, Pudipatla post, Tirupathi, Andhra Pradesh 517505, India

plus web supplement of 1 page

Fracture repair in domestic animals is a routine procedure but the same in wild animals is rare. Though fractures do occur in wild animals, lack of personnel to capture, facilities to perform surgery, complications during anaesthesia, problems in postoperative management prevent anybody to make an attempt. In this present study a case report of bone plating for tibial fracture repair in a wild Black Buck *Antelope cervicapra* is discussed with detailed anaesthetic and surgical procedure.

Case history: A female Black Buck aged around 12 years was presented to the college hospital with history of limping in the left hind limb. The zoo veterinarian had treated the case with analgesics and anti-inflammatory drugs. Temperature, pulse and respiration were within the normal range. Palpation of mid shaft of tibia revealed crepitation and the animal was also evincing pain. No external injury was noticed. Plain radiography revealed complete fracture of tibial mid shaft (Image 1^w). It was decided to perform internal stabilization under general anaesthesia.

Anaesthesia and surgical procedure: The animal was prepared for surgery; food and water withheld for 12 hours. Preoperatively, 500mg of metronidazole, 5% dextrose normal saline and ciprofloxacin were injected i/v. The animal was anaesthetized using combination of xylazine hydrochloride (40mg) and ketamine hydrochloride (150mg) given i/m. The animal was positioned in lateral recumbancy. The medial aspect of tibia was prepared for aseptic surgery. A 6cm cutaneous incision was made and separated fascia and muscles. The proximal and distal fractured segments of the bone were identified and the site irrigated with normal saline solution to remove debris and blood clots.

^w See Images 1 & 2 in the web supplement at www.zoosprint.org

Using extension and counter extension the two segments were reduced to normal alignment. Eight holed Shearman's stainless steel bone plate was kept inside and 3mm cortical screws were used to stabilize the bone plate to the fractured segments (Image 2^w). Also circlage wire was applied to prevent loosening of the plate. The muscles were sutured using no.1 chromic catgut. Sub cuticular sutures were applied to prevent dead space. Cutaneous incision was closed using no.1 braided silk. Tincture benzoin seal was applied to the suture line. Plaster of paris cast was applied to the limb for external immobilization leaving an adequate window for dressing of surgical wound. The animal received metronidazole (500mg), ciprofloxacin (500mg), 5% dextrose normal saline (1000ml) i/v for five post operative days along with dressing of wound.

Xylazine and ketamine provided satisfactory anaesthesia for the entire surgical procedure without any complications (Lumb & Jones, 1996). Similar anaesthetic protocols were also reported by Jalynski (2003) for treating fractures in sheep. Bone plating with cortical screws provided effective stabilization of fractured segments. External immobilization with plaster of paris cast prevented movement of segments and allowed regular dressing of the surgical wound. Pre-, intra- and postoperative use of antibiotics and fluids provided aseptic conditions of site without any problems.

References

- Jalynski, M. (2003). The use of an external fixator MACZEK to treat fractures in sheep. *Indian Veterinary Journal* 80: 642-646.
Lumb, W.V. & E.W. Jones (1996). Anaesthesia of wild and laboratory animals, pp.698-701. In: *Veterinary Anaesthesia*. 3rd edition. Williams and Williams Maryland USA.

Acknowledgement: Authors acknowledge the necessary permission from the Associated Dean, Dr. V. Haragopal, Professor of Surgery College of Veterinary Science; A. Partha Sarathy, In-charge Curator, S.V. Zoo Park; sergeant and the animal attendants for their help during this work.



VET BRIEF

ZOOS' PRINT JOURNAL 22(6): 2736-2737

Occurrence of *Babesia* infection in Leopard *Panthera pardus* at Nagpur

B.S. Baviskar¹, P.J. Gawande², D.K. Maske³, A.K. Jayraw⁴, S.S. Bawaskar⁵ and A.G. Bhandarkar⁶

^{1,2,3,4,6} Department of Parasitology, Nagpur Veterinary College, M.A.F.S.U., Seminary Hills, Nagpur, Maharashtra 440006, India
⁵ Maharajbagh Zoo, College of Agriculture, Nagpur, Maharashtra, India
Email: ¹ drbaharbaviskar@rediffmail.com; ² drpriyagawande@gmail.com; ⁴ jayrawanant1@rediffmail.com

plus web supplement of 1 page

Babesiosis is a tick-borne haemoprotozoan disease caused by *Babesia* sp. and characterized by pyrexia, haemolytic anaemia, haemoglobinuria, jaundice and death. In India, babesiosis is widespread and recognized as one of the serious problem of major economic significance in domestic animals, although, sufficient information is not available on incidence of the disease in wild animals (Arora, 1994). Hence, these cases of babesiosis in Leopard *Panthera pardus* from Maharajbagh Zoo, Nagpur, Maharashtra are reported here.

Two 15-year old male leopards died at Maharajbagh Zoo, College of Agriculture, Nagpur with a history of anorexia, dehydration, convulsions and lumbar pain. Postmortem examination was performed; heart blood smears were prepared, stained with Leishman's stain and

^w See Image 1 in the web supplement at www.zoosprint.org